

- A magnetic field of flux density 1.0Wbm^{-2} acts normal to a 80 turn coil of 0.01m^2 area. If this coil is removed from the field in 0.2 second, the emf induced in it is

(A) 5V (B) 4V (C) 8V (D) 0.8V
- An alternative current is given by $i = i_1 \sin \omega t + i_2 \cos \omega t$. The r.m.s. current is given by

(A) $\sqrt{\frac{i_1^2 + i_2^2}{2}}$ (B) $\frac{i_1 + i_2}{\sqrt{2}}$ (C) $\frac{i_1 i_2}{\sqrt{2}}$ (D) $\sqrt{\frac{i_1^2 + i_2^2}{2}}$
- Which of the following statements proves that Earth has a magnetic field?

(A) A large quantity of iron- ore is found in the Earth
 (B) The intensity of cosmic rays stream of charged particles is more at the pole than at the equator.
 (C) Earth is planet rotating about the North South axis.
 (D) Earth is surrounded by ionosphere.
- A long solenoid has 500 turns, when a current of 2A is passed through it, the resulting magnetic flux linked with each turn of the solenoid is $4 \times 10^{-3}\text{Wb}$, then self induction of the solenoid is

(A) 1.0 henry (B) 4.0 henry (C) 2.5 henry (D) 2.0 henry
- A fully charged capacitor 'C' with initial charge 'q₀' is connected to a coil self inductance 'L' at t=0. The time at which the energy is stored equally between the electric and the magnetic field is

(A) $\frac{\pi}{4}\sqrt{LC}$ (B) $2\pi\sqrt{LC}$ (C) \sqrt{LC} (D) $\pi\sqrt{LC}$
- The power of a equi- concave lens is -4.5D and is made od a material of R.I. 1.6, the radii of curvature of the lens is


(A) 115.44cm (B) -26.6cm (C) $+36.6\text{cm}$ (D) -2.66cm
- A ray of light passes through an equilateral glass prism in such a manner that the angle of incidence is equal to the angle of emergence and each of these angles is equal to $\frac{3}{4}$ of the angle of the prism. The angle of deviation is

(A) 30° (B) 45° (C) 39° (D) 20°
- A convex lens of focal length 'f' is placed somewhere in between an object and a screen. The distance between the object and the screen is 'x'. If the numerical value of the magnification produced by the lens is 'm', then the focal length of the lens is


(A) $\frac{(m-1)^2 x}{m}$ (B) $\frac{mx}{(m+1)^2}$ (C) $\frac{mx}{(m-1)^2}$ (D) $\frac{(m+1)^2 x}{m}$
- A series resonant ac circuit contains a capacitance 10^{-6}F and an inductor of 10^{-4}H . The frequency of electrical oscillation will be

(A) $\frac{10}{2\pi}\text{Hz}$ (B) 10^5Hz (C) 10Hz (D) $\frac{10^5}{2\pi}\text{Hz}$

10. In a series LCR circuit $R = 300\Omega$, $L = 0.9H$, $C = 2.0\mu F$ and $\omega = 1000\text{ rad s}^{-1}$, then impedance of the circuit is
 (A) 400Ω (B) 1300Ω (C) 900Ω (D) 500Ω
11. Which of the following radiations of electromagnetic waves has the highest wave length?
 (A) Microwaves (B) X- rays (C) UV- rays (D) IR- rays
12. The fringe width for red colour as compared to that for violet colour is approximately
 (A) 8 times (B) 3 times (C) Double (D) 4 times
13. In case of Fraunhofer diffraction at a single slit the diffraction pattern on the screen is correct for which of the following statements?
 (A) Central bright band having dark bands on either side.
 (B) Central dark band having alternate dark and bright bands of decreasing intensity on either side.
 (C) Central bright band having alternate dark and bright bands of decreasing intensity on either side
 (D) Central dark band having uniform brightness on either side.
14. When a Compact Disc (CD) is illuminated by small source of white light coloured bands are observed. This is due to
 (A) Reflection (B) Scattering (C) Diffraction (D) Interference
15. Consider a glass slab which is silvered at one side and the other side is transparent. Given the refractive index of the glass slab to be 1.5. If a ray of light is incident at an angle of 45° on the transparent side, the deviation of the ray of light from its initial path, when it comes out of the slab is
 (A) 45° (B) 90° (C) 180° (D) 120°
16. Focal length of a convex lens will be maximum for
 (A) Red light (B) Blue light (C) Yellow light (D) Green light
17. For light diverging from a finite point source
 (A) The intensity at the wave front does not depend on the distance.
 (B) The wave front is cylindrical
 (C) The intensity decreases in proportion to the distance squared.
 (D) The wave front is parabolic.
18. The radius of hydrogen atom in the ground state is 0.53 \AA . After collision with an electrons, it is found to have a radius of 2.12 \AA , the principal quantum number 'n' of the final state of the atom is
 (A) $n = 4$ (B) $n = 1$ (C) $n = 2$ (D) $n = 3$
19. In accordance with the Bohr's model, the quantum number that characterises the Earth's revolution around the Sun in an orbit of radius $1.5 \times 10^{11}\text{ m}$ with orbital speed $3 \times 10^4\text{ m s}^{-1}$ is [given mass of Earth = $6 \times 10^{24}\text{ kg}$]
 (A) 2.57×10^{74} (B) 5.98×10^{86} (C) 2.57×10^{38} (D) 8.57×10^{64}
20. If an electron is revolving in its Bohr orbit having Bohr radius of 0.529 \AA , then the radius of third orbit is
 (A) 5125 nm (B) 4234 nm (C) 4496 \AA (D) 4.761 \AA

21. Binding energy of a Nitrogen nucleus ${}_{7}^{14}\text{N}$, given $m[{}_{7}^{14}\text{N}] = 14.00307u$
- (A) 78 MeV (B) 104.7 MeV (C) 85 MeV (D) 206.5 MeV
22. In a photo electric experiment, if both the intensity and frequency of the incident light are doubled, then the saturation photo electric current
- (A) becomes four times (B) remains constant
(C) is halved (D) is doubled
23. The kinetic energy of the photoelectrons increases by 0.52 eV when the wavelength of incident light is changed from 500 nm to another wavelength which is approximately
- (A) 1000 nm (B) 700 nm (C) 400 nm (D) 1250 nm
24. The de-Broglie wavelength of a particle of kinetic energy 'K' is λ ; the wavelength of the particle, if its kinetic energy is $\frac{K}{4}$ is
- (A) 4λ (B) λ (C) 2λ (D) $\frac{\lambda}{2}$
25. The forbidden energy gap of germanium crystal at '0' K is
- (A) 6.57 eV (B) 0.071eV (C) 0.71 eV (D) 2.57 eV
26. Which logic gate is represented by the following combination of logic gates?
- 
- (A) NOR (B) OR (C) NAND (D) AND
27. A metallic rod of mass per unit length 0.5kg m^{-1} is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. A magnetic field of strength 0.25 T is acting on it in the vertical direction. When a current 'I' is flowing through it, the rod is not allowed to slide down. The quantity of current required to keep the rod stationary is
- (A) 11.32 A (B) 7.14 A (C) 5.98 A (D) 14.76 A
28. A nuclear reactor delivers a power of 10^9 W , the amount of fuel consumed by the reactor in one hour is
- (A) 0.96 g (B) 0.04 g (C) 0.08 g (D) 0.72 g
29. Which of the following radiations is deflected by electric field?
- (A) α - particles (B) X-rays (C) Neutrons (D) γ - rays
30. The resistivity of a semiconductor at room temperature is in between
- (A) 10^{10} to $10^{12}\Omega\text{ cm}$ (B) 10^{-2} to $10^{-5}\Omega\text{ cm}$ (C) 10^{-3} to $10^6\Omega\text{ cm}$ (D) 10^6 to $10^8\Omega\text{ cm}$
31. A Car is moving on a circular horizontal track of radius 10 m with a constant speed of 10 ms^{-1} . A bob is suspended from the roof of the car by a light wire of length 1.0 m. The angle made by the vertical is (in radian)
- (A) $\frac{\pi}{3}$ (B) $\frac{\pi}{6}$ (C) $\frac{\pi}{4}$ (D) 0

32. Two masses of 5 kg and 3 kg are suspended with the help of massless inextensible strings as shown in figure. When system is going upwards with acceleration 2 ms^{-2} , the value of T_1 is (use $g = 9.8 \text{ ms}^{-2}$)



- (A) 59 N (B) 94.4 N (C) 35.4 N (D) 23.6 N
33. The Vernier scale of a travelling microscope has 50 division which coincides with 49 main scale divisions. If each main scale division is 0.5 mm, then the least count of the microscope is
- (A) 0.5 cm (B) 0.01 cm (C) 0.5 mm (D) 0.01 mm
34. The displacement ' x ' (in metre) of particle of mass ' m ' (in kg) moving in one dimension under the action of a force, is related to time ' t ' (in sec.) by, $t = \sqrt{x} + 3$. The displacement of the particle when its velocity is zero, will be
- (A) $2m$ (B) $4m$ (C) $0m$ (D) $6m$
35. Two objects are projected at an angle θ° and $(90 - \theta)^\circ$, to the horizontal with the same speed. The ratio of their maximum vertical height is
- (A) $\tan^2 \theta : 1$ (B) 1:1 (C) $\tan \theta : 1$ (D) $1 : \tan \theta$
36. A metallic rod breaks when strain produced is 0.2%. The Young's modulus the materials of the rod is $7 \times 10^9 \text{ Nm}^{-2}$. The area of cross section to support a load of 10^4 N is
- (A) $7.1 \times 10^{-2} \text{ m}^2$ (B) $7.1 \times 10^{-8} \text{ m}^2$ (C) $7.1 \times 10^{-6} \text{ m}^2$ (D) $7.1 \times 10^{-4} \text{ m}^2$
37. A tiny spherical oil drop carrying a net charge q is balanced in still air, with a vertical uniform electric field of strength $\frac{81}{7} \pi \times 10^5 \text{ V/m}$. When the field is switched off, the drop is observed to fall with terminal velocity $2 \times 10^{-3} \text{ ms}^{-1}$. Here $g = 9.8 \text{ ms}^{-2}$, viscosity of air is $1.8 \times 10^{-5} \text{ N s/m}^2$ and the density of oil is 900 kg m^{-3} . The magnitude of ' q ' is
- (A) $3.2 \times 10^{-19} \text{ C}$ (B) $0.8 \times 10^{-19} \text{ C}$ (C) $8 \times 10^{-19} \text{ C}$ (D) $1.6 \times 10^{-19} \text{ C}$
38. "Heat cannot be itself flow from a body at lower temperature to a body at higher temperature". This statement corresponds to
- (A) First law of thermodynamics (B) Second law of Thermodynamics
(C) Conservation of momentum (D) Conservation of mass
39. A smooth chain of length 2 m is kept on a table such that its length of 60 cm hangs freely from the edge of the table. The total mass of the chain is 4 kg. The work done in pulling the entire chain on the table is, (Take $g = 10 \text{ ms}^{-2}$)
- (A) 2.0 J (B) 12.9 J (C) 6.3 J (D) 3.6 J

40. The angular speed of a motor wheel is increased from 1200 rpm to 3120 rpm in 16 seconds.
The angular acceleration of the motor wheel is
(A) $8\pi \text{ rad s}^{-2}$ (B) $2\pi \text{ rad s}^{-2}$ (C) $4\pi \text{ rad s}^{-2}$ (D) $6\pi \text{ rad s}^{-2}$
41. The centre of mass of an extended body on the surface of the earth and its centre of gravity
(A) centre of mass coincides with the centre of gravity of a body if the size of the body is negligible as compared to the size (or radius) of the earth
(B) are always at the same point for any size of the body.
(C) Are always at the same point only for spherical bodies.
(D) Can never be at the same point.
42. An electric dipole with dipole moment $4 \times 10^{-9} \text{ Cm}$ is aligned at 30° with the direction of a uniform electric field of magnitude $5 \times 10^4 \text{ NC}^{-1}$, the magnitude of the torque acting on the dipole is
(A) $10 \times 10^{-3} \text{ Nm}$ (B) 10^{-4} Nm (C) $\sqrt{3} \times 10^{-4} \text{ Nm}$ (D) 10^{-5} Nm
43. A charged particle of mass ' m ' and charge ' q ' is released from rest in an uniform electric field \vec{E} . Neglecting the effect of gravity, the kinetic energy of the charged particle after ' t ' second is
(A) $\frac{E^2 q^2 t^2}{2m}$ (B) $\frac{2E^2 t^2}{mq}$ (C) $\frac{Eq^2 m}{2t^2}$ (D) $\frac{Eqm}{t}$
44. The electric field and the potential of an electric dipole vary with distance r as
(A) $\frac{1}{r^3}$ and $\frac{1}{r^2}$ (B) $\frac{1}{r}$ and $\frac{1}{r^2}$ (C) $\frac{1}{r^2}$ and $\frac{1}{r}$ (D) $\frac{1}{r^2}$ and $\frac{1}{r^3}$
45. The displacement of a particle executing SHM is given by $X = 3 \sin \left[2\pi t + \frac{\pi}{4} \right]$ where ' x ' is in metres and ' t ' is in seconds. The amplitude and maximum speed of the particle is
(A) 3 m, $8\pi \text{ ms}^{-1}$ (B) 3 m, $2\pi \text{ ms}^{-1}$ (C) 3 m, $4\pi \text{ ms}^{-1}$ (D) 3 m, $6\pi \text{ ms}^{-1}$
46. Electrical as well as gravitational effects can be thought to be caused by fields. Which of the following is true for an electrical or gravitational field?
(A) There is no way to verify the existence of force field since it is just a concept
(B) The field concept is often used to describe contact forces.
(C) Gravitational or Electric fields does not exist in the space around an object
(D) Fields are useful for understanding forces acting through a distance.
47. Four charges $+q, +2q, +q$ and $-2q$ are placed at the corners of a square $ABCD$ respectively.
The force on a unit positive charge kept at the centre ' O ' is
(A) perpendicular to AD (B) zero
(C) along the diagonal BD (D) along the diagonal AC
48. Wire bound resistors are made by
(A) winding the wires of an alloy of manganin, constantan, nichrome
(B) winding the wires of an alloy of $\text{Cu}, \text{Al}, \text{Ag}$
(C) winding the wires of an alloy of $\text{Si}, \text{Tu}, \text{Fe}$
(D) winding the wires of an alloy of $\text{Ge}, \text{Au}, \text{Ga}$

49. Ten identical cells each of potential ' E ' and internal resistance ' r ' are connected in series to form a closed circuit. An ideal voltmeter connected across three cells, will read
- (A) $7E$ (B) $10E$ (C) $3E$ (D) $13E$
50. In an atom electrons revolve around the nucleus along a path of radius 0.72 \AA making 9.4×10^{18} revolutions per second. The equivalent current is $-\left[\text{given } e = 1.6 \times 10^{-19} \text{ C}\right]$
- (A) 1.8 A (B) 1.2 A (C) 1.5 A (D) 1.4 A
51. When a metal conductor connected to left gap of a meter bridge is heated, the balancing point
- (A) shifts to the centre (B) shifts towards right
(C) shifts towards left (D) remains unchanged
52. Two tiny spheres carrying charges $1.8 \mu\text{C}$ and $2.8 \mu\text{C}$ are located at 40 cm apart. The potential at the mid-point of the line joining the two charges is
- (A) $3.6 \times 10^5 \text{ V}$ (B) $3.8 \times 10^4 \text{ V}$ (C) $21 \times 10^5 \text{ V}$ (D) $4.3 \times 10^4 \text{ V}$
53. A parallel plate capacitor is charged by connecting a 2 V battery across it. It is then disconnected from the battery and a glass slab is introduced between plates. Which of the following pairs of quantities decrease?
- (A) Capacitance and charge (B) Charge and potential difference
(C) Potential difference and energy stored (D) Energy stores and capacitance.
54. A charged particle is moving in an electric field of $3 \times 10^{-10} \text{ V m}^{-1}$ with mobility $2.5 \times 10^6 \text{ m}^2/\text{V/s}$, its drift velocity is
- (A) $1.2 \times 10^{-4} \text{ ms}^{-1}$ (B) $7.5 \times 10^{-4} \text{ ms}^{-1}$ (C) $8.33 \times 10^{-4} \text{ ms}^{-1}$ (D) $2.5 \times 10^4 \text{ ms}^{-1}$
55. A solenoid of length 50 cm having 100 turns carries a current of 2.5 A . The magnetic field at one end of the solenoid is
- (A) $9.42 \times 10^{-4} \text{ T}$ (B) $3.14 \times 10^{-4} \text{ T}$ (C) $6.285 \times 10^{-4} \text{ T}$ (D) $1.57 \times 10^{-4} \text{ T}$
56. A galvanometer of resistance 50Ω is connected to a battery of 3 V along with a resistance 2950Ω in series. A full scale deflection of 30 divisions is obtained in the galvanometer. In order to reduce this deflection to 20 divisions, the resistance in series should be
- (A) 4450Ω (B) 6050Ω (C) 5550Ω (D) 5050Ω
57. A circular coil of wire of radius ' r ' has ' n ' turns and carries a current ' I '. The magnetic induction ' B ' at a point on the axis of the coil at a distance $\sqrt{3} r$ from its centre is
- (A) $\frac{\mu_0 n I}{4r}$ (B) $\frac{\mu_0 n I}{32r}$ (C) $\frac{\mu_0 n I}{8r}$ (D) $\frac{\mu_0 n I}{16r}$
58. If voltage across a bulb rated $220 \text{ V}, 100 \text{ W}$ drops by 2.5% of its rated value, the percentage of the rated value by which the power would decrease is
- (A) 10% (B) 20% (C) 2.5% (D) 5%

