

1. The value of $\cot^{-1} \left[\frac{\sqrt{1-\sin x} + \sqrt{1+\sin x}}{\sqrt{1-\sin x} - \sqrt{1+\sin x}} \right]$ where $x \in \left(0, \frac{\pi}{4}\right)$ is

(A) $\pi - \frac{x}{3}$

(B) $\frac{x}{2}$

(C) $\pi - \frac{x}{2}$

(D) $\frac{x}{2} - \pi$

Sol: $\cot^{-1} \left[\frac{\sqrt{1-\sin x} + \sqrt{1+\sin x}}{\sqrt{1-\sin x} - \sqrt{1+\sin x}} \right], x \in \left(0, \frac{\pi}{4}\right)$

$$\sqrt{1+\sin x} = \cos \frac{x}{2} + \sin \frac{x}{2}$$

$$\sqrt{1-\sin x} = \sqrt{\left(\cos \frac{x}{2} - \sin \frac{x}{2}\right)^2} = \cos \frac{x}{2} - \sin \frac{x}{2}$$

$$\text{G.E.} = \cot^{-1} \left[\frac{\cos \frac{x}{2} + \sin \frac{x}{2} + \cos \frac{x}{2} - \sin \frac{x}{2}}{\cos \frac{x}{2} + \sin \frac{x}{2} - \left(\cos \frac{x}{2} - \sin \frac{x}{2}\right)} \right]$$

$$= \cot^{-1} \left[\cot \frac{x}{2} \right] = \frac{x}{2}$$

Ans: (B)

2. If $x \begin{bmatrix} 3 \\ 2 \end{bmatrix} + y \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 15 \\ 5 \end{bmatrix}$ then the value of x and y are

(A) $x = -4, y = -3$

(B) $x = 4, y = 3$

(C) $x = -4, y = 3$

(D) $x = 4, y = -3$

Sol:

$$3x + y = 15$$

$$2x - y = 5$$

$$5x = 20$$

$$x = 4$$

$$3(4) + y = 15$$

$$y = 15 - 12$$

$$y = 3$$

Ans: (B)

3. If A and B are two matrices such that $AB = B$ and $BA = A$ then $A^2 + B^2 =$

(A) AB

(B) $A+B$

(C) $2BA$

(D) $2AB$

Sol: $A^2 + B^2 = A \cdot A + B \cdot B = A(BA) + B(AB) = (AB)A + (BA)B$

$$= BA + AB = A + B$$

Ans: (B)

4. If $A = \begin{bmatrix} 2-k & 2 \\ 1 & 3-k \end{bmatrix}$ is singular matrix, then the value of $5k - k^2$ is equal to

(A) -4

(B) 4

(C) 6

(D) -6

Sol: $(2-k)(3-k) - 2 = 0$

$$6 - 5k + k^2 - 2 = 0$$

$$4 = 5k - k^2$$

Ans: (B)

5. The area of a triangle with vertices $(-3,0)$, $(3,0)$ and $(0,k)$ is 9 sq. units, the value of k is

(A) 6

(B) 9

(C) 3

(D) -9

Sol: $9 = \frac{1}{2} \begin{vmatrix} -3 & 0 & 1 \\ 3 & 0 & 1 \\ 0 & k & 1 \end{vmatrix}$

$$\pm 18 = -3(-k) + 1(3k)$$

$$\pm 18 = 6k \Rightarrow k = \pm 3$$

Ans: (C)

6. If $\Delta = \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix}$ and $\Delta_1 = \begin{vmatrix} 1 & 1 & 1 \\ bc & ca & ab \\ a & b & c \end{vmatrix}$ then

(A) $\Delta_1 \neq \Delta$

(B) $\Delta_1 = \Delta$

(C) $\Delta_1 = -\Delta$

(D) $\Delta_1 = 3\Delta$

Sol: $D = \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix}$

$$R_2 \rightarrow R_1 - R_2 \text{ and } R_3 \rightarrow R_3 - R_1$$

$$\Delta = \begin{vmatrix} 1 & a & a^2 \\ 0 & a-b & (a+b)(a-b) \\ 0 & c-a & (c+a)(c-a) \end{vmatrix}$$

$$\Delta = (a-b)(c-b)(c-a)$$

$$\Delta_1 = \begin{vmatrix} 1 & 1 & 1 \\ bc & ca & ab \\ a & b & c \end{vmatrix}$$

$$C_2 \rightarrow C_2 - C_1, C_3 \rightarrow C_3 - C_1$$

$$\Delta_1 = \begin{vmatrix} 1 & 0 & 0 \\ bc & c(a-b) & b(a-c) \\ a & b-a & c-a \end{vmatrix}$$

$$\Delta_1 = (a-b)(c-b)(c-a)$$

Ans: (B)

7. If $\sin^{-1}\left(\frac{2a}{1+a^2}\right) + \cos^{-1}\left(\frac{1-a^2}{1+a^2}\right) = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$ where $a, x \in (0,1)$ then the value of x is
- (A) $\frac{2a}{1+a^2}$ (B) 0 (C) $\frac{2a}{1-a^2}$ (D) $\frac{a}{2}$

Sol: $\sin^{-1}\left(\frac{2a}{1+a^2}\right) + \cos^{-1}\left(\frac{1-a^2}{1+a^2}\right) = \tan^{-1}\left(\frac{2x}{1-x^2}\right), a, x \in (0,1)$

$\Rightarrow 2 \tan^{-1} a + 2 \tan^{-1} a = 2 \tan^{-1} x$

$\Rightarrow 2 \tan^{-1} a = \tan^{-1} x$

$\Rightarrow \tan^{-1}\left(\frac{2a}{1-a^2}\right) = \tan^{-1} x \quad \therefore x = \frac{2a}{1-a^2}$

Ans: (C)

8. If $u = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$ and $v = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$ then $\frac{du}{dv}$ is
- (A) $\frac{1-x^2}{1+x^2}$ (B) $\frac{1}{2}$ (C) 1 (D) 2

Sol: $u = \sin^{-1}\left(\frac{2x}{1+x^2}\right) \quad v = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$

$u = 2 \tan^{-1} x \quad v = 2 \tan^{-1} x$

$\frac{du}{dx} = \frac{2}{1+x^2} \quad \frac{dv}{dx} = \frac{2}{1+x^2}$


$\therefore \frac{du}{dv} = 1$

Ans: (C)

9. The function $f(x) = \cot x$ is discontinuous on every point of the set

- (A) $\left\{x = (2n+1)\frac{\pi}{2}; n \in \mathbb{Z}\right\}$ (B) $\{x = n\pi; n \in \mathbb{Z}\}$
- (C) $\left\{x = \frac{n\pi}{2}; n \in \mathbb{Z}\right\}$ (D) $\{x = 2n\pi; n \in \mathbb{Z}\}$

Sol:



Graph of $\cot x$

$$\cot x = \frac{\cos x}{\sin x} \text{ is undefined when } \sin x = 0$$

$\therefore \cot x$ is undefined at $x = n\pi, n \in \mathbb{Z}$

Ans: (B)

10. If the function is $f(x) = \frac{1}{x+2}$, then the point of discontinuity of the composite function $y = f(f(x))$ is

- (A) $\frac{2}{5}$ (B) $\frac{-5}{2}$ (C) $\frac{1}{2}$ (D) $\frac{5}{2}$

Sol: $f(x) = \frac{1}{x+2}$ is undefined at $x = -2$

$$f(f(x)) = \frac{1}{f(x)+2} = \frac{1}{\frac{1}{x+2}+2} = \frac{x+2}{2x+5} \text{ is undefined at } x = -\frac{5}{2}$$

Ans: (B)

11. If $y = a \sin x + b \cos x$, then $y^2 + \left(\frac{dy}{dx}\right)^2$ is a

- (A) function of x and y (B) function of x
 (C) constant (D) function of y

Sol: $y = a \sin x + b \cos x$

$$\begin{aligned} y^2 + \left(\frac{dy}{dx}\right)^2 &= (a \sin x + b \cos x)^2 + (a \cos x - b \sin x)^2 \\ &= a^2 \sin^2 x + b^2 \cos^2 x + 2ab \sin x \cos x + a^2 \cos^2 x + b^2 \sin^2 x - 2ab \sin x \cos x \\ &= a^2 + b^2 \text{ is constant} \end{aligned}$$

Ans: (C)

12. If $f(x) = 1 + nx + \frac{n(n-1)}{2}x^2 + \frac{n(n-1)(n-2)}{6}x^3 + \dots + x^n$ then $f''(1) =$

- (A) $n(n-1)2^n$ (B) $(n-1)2^{n-1}$ (C) 2^{n-1} (D) $n(n-1)2^{n-2}$

Sol: $f(x) = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + x^n$

$$= (1+x)^n$$

$$f'(x) = n(1+x)^{n-1}$$

$$f''(x) = n(n-1)(1+x)^{n-2}$$

$$f''(1) = n(n-1) \cdot 2^{n-2}$$

Ans: (D)

13. If $A = \begin{bmatrix} 1 & \tan\alpha/2 \\ -\tan\alpha/2 & 1 \end{bmatrix}$ and $AB = I$ then $B =$

- (A) $\cos^2\alpha/2 \cdot I$ (B) $\cos^2\alpha/2 \cdot A^T$ (C) $\sin^2\alpha/2 \cdot A$ (D) $\cos^2\alpha/2 \cdot A$

Sol: $B = A^{-1}$

$$A = \begin{bmatrix} 1 & \tan\frac{\alpha}{2} \\ -\tan\frac{\alpha}{2} & 1 \end{bmatrix}$$

$$|A| = 1 + \tan^2\frac{\alpha}{2}$$

$$= \sec^2\frac{\alpha}{2}$$

$$A^{-1} = \frac{1}{\sec^2\frac{\alpha}{2}} \begin{bmatrix} 1 & -\tan\frac{\alpha}{2} \\ \tan\frac{\alpha}{2} & 1 \end{bmatrix} = \cos^2\frac{\alpha}{2} A^T$$

Ans: (B)

14. A circular plate of radius 5cm is heated. Due to expansion, its radius increases at the rate of 0.05cm/sec. The rate at which its area is increasing when the radius is 5.2cm is

- (A) $5.05\pi\text{cm}^2/\text{sec}$ (B) $5.2\pi\text{cm}^2/\text{sec}$ (C) $0.52\pi\text{cm}^2/\text{sec}$ (D) $27.4\pi\text{cm}^2/\text{sec}$


Sol: $\frac{dr}{dt} = 0.05 \text{ cm/sec}$

$$A = \pi r^2$$

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

$$\left(\frac{dA}{dt}\right) = 2\pi(5.2)(0.05) = 0.52\pi$$

Ans: (C)



15. The distance 's' in meters travelled by a particle in 't' seconds is given by $s = \frac{2t^3}{3} - 18t + \frac{5}{3}$. The acceleration when the particle comes to rest is

- (A) $12\text{m}^2/\text{sec}$. (B) $3\text{m}^2/\text{sec}$. (C) $18\text{m}^2/\text{sec}$. (D) $10\text{m}^2/\text{sec}$.

Sol: $s = \frac{2t^3}{3} - 18t + \frac{5}{3}$

$$v = \frac{ds}{dt} = 2t^2 - 18$$

$$a = \frac{dv}{dt} = 4t \quad v = 0 \Rightarrow t = 3$$

$$\therefore a = 12$$

Ans: (A)

16. A particle moves along the curve $\frac{x^2}{16} + \frac{y^2}{4} = 1$. When the rate of change of abscissa is 4 times that of its ordinate, then the quadrant in which the particle lies is

(A) III or IV (B) I or III (C) II or III (D) II or IV

Sol: $\frac{x^2}{16} + \frac{y^2}{4} = 1$

$$\frac{2x}{16} \frac{dx}{dt} + \frac{2y}{4} \frac{dy}{dt} = 0$$

$$\frac{x}{8} \frac{dx}{dt} + \frac{y}{2} \frac{dy}{dt} = 0$$

$$\Rightarrow \frac{x}{8} \left(4 \frac{dy}{dt} \right) + \frac{y}{2} \frac{dy}{dt} = 0$$

$$\Rightarrow \frac{dy}{dt} \left(\frac{x}{2} + \frac{y}{2} \right) = 0$$

$$\Rightarrow x = -y$$

$$\Rightarrow (x, y) \text{ lie in II or IV}$$

Ans: (D)

17. An enemy fighter jet is flying along the curve given by $y = x^2 + 2$. A soldier is placed at (3, 2) wants to shoot down the jet when it is nearest to him. Then the nearest distance is

(A) 2 units (B) $\sqrt{3}$ units (C) $\sqrt{5}$ units (D) $\sqrt{6}$ units

Sol: $y = x^2 + 2 \quad \therefore p(t, t^2 + 2)$

A(3, 2)

Slope of AP = $\frac{t^2}{t-3}$

$y = x^2 + 2 \Rightarrow y' = 2x$

Slope of normal = Slope of AP

$$\frac{-1}{2t} = \frac{t^2}{t-3}$$

$$2t^3 = -t + 3$$

$$2t^3 = -t + 3$$

$$2t^3 + t - 3 = 0$$

$$t = 0 \quad \therefore p = (1, 3)$$

$$\therefore AP = \sqrt{4+1} = \sqrt{5}$$

Ans: (C)

18. $\int_2^8 \frac{5\sqrt{10-x}}{5\sqrt{x} + 5\sqrt{10-x}} dx =$

- (A) 4 (B) 5 (C) 3 (D) 6

Sol: $I = \int_2^8 \frac{5\sqrt{10-x}}{5\sqrt{x} + 5\sqrt{10-x}} dx \quad \dots (1)$

$I = \int_2^8 \frac{5\sqrt{x}}{5\sqrt{10-x} + 5\sqrt{x}} dx \quad \dots (2)$

$2I = \int_2^8 1 dx = x \Big|_2^8 = 8 - 2 = 6$

$I = 3$

Ans: (C)

19. $\int \sqrt{\operatorname{cosec} x - \sin x} dx =$

- (A) $2\sqrt{\sin x} + C$ (B) $\sqrt{\sin x} + C$ (C) $\frac{2}{\sqrt{\sin x}} + C$ (D) $\frac{\sqrt{\sin x}}{2} + C$

Sol: $\int \sqrt{\operatorname{cosec} x - \sin x} dx = \int \sqrt{\frac{1}{\sin x} - \sin x} dx$

$= \int \sqrt{\frac{1 - \sin^2 x}{\sin x}} dx$

$\int \frac{\cos x}{\sqrt{\sin x}} dx \quad \sin x = t, \cos x dx = dt$

$= \int \frac{1}{\sqrt{t}} dt$

$= 2\sqrt{\sin x} + C$

Ans: (A)

20. If $f(x)$ and $g(x)$ are two functions with $g(x) = x - \frac{1}{x}$ and $f \circ g(x) = x^3 - \frac{1}{x^3}$ then $f'(x) =$

- (A) $x^2 - \frac{1}{x^2}$ (B) $3x^2 + 3$ (C) $1 - \frac{1}{x^2}$ (D) $3x^2 + \frac{3}{x^4}$

Sol: $g(x) = x - \frac{1}{x} \Rightarrow (f \circ g)(x) = x^3 - \frac{1}{x^3}$

$f(g(x)) = x^3 - \frac{1}{x^3} = \left(x - \frac{1}{x}\right)^3 + 3\left(x - \frac{1}{x}\right)$

$\Rightarrow f(g(x)) = (g(x))^3 + 3g(x)$

$\Rightarrow f(x) = x^3 + 3x \Rightarrow f'(x) = 3x^2 + 3$

Ans: (B)

21. $\int \frac{1}{1+3\sin^2 x+8\cos^2 x} dx =$

- (A) $\frac{1}{6} \tan^{-1}\left(\frac{2\tan x}{3}\right) + C$ (B) $\frac{1}{6} \tan^{-1}(2\tan x) + C$ (C) $6\tan^{-1}\left(\frac{2\tan x}{3}\right) + C$ (D) $\tan^{-1}\left(\frac{2\tan x}{3}\right) + C$

Sol: $\int \frac{1}{1+3\sin^2 x+8\cos^2 x} dx$

$$= \int \frac{\sec^2 x}{\sec^2 x + 3\tan^2 x + 8} dx$$

$$= \int \frac{\sec^2 x dx}{9+4\tan^2 x}$$

$$= \int \frac{dt}{3^2 + (2t)^2}$$

$\tan x = t, \sec^2 x dx = dt$

$$= \frac{1}{3} \tan^{-1}\left(\frac{2t}{3}\right) \cdot \frac{1}{2}$$

$$= \frac{1}{6} \tan^{-1}\left(\frac{2\tan x}{3}\right)$$

Ans: (A)

22. $\int_{-2}^0 (x^3 + 3x^2 + 3x + 3 + (x+1)\cos(x+1)) dx =$

- (A) 4 (B) 0 (C) 1 (D) 3

Sol: $\int_{-2}^0 ((x+1)^3 + 2 + (x+1)\cos(x+1)) dx$

$$\int_{-1}^1 (x^3 + 2 + x \cos x) dx$$

$$2x \Big|_{-1}^1 = 2(1+1) = 4$$

Ans: (A)

23. $\int_0^{\pi} \frac{x \tan x}{\sec x \cdot \operatorname{cosec} x} dx =$

- (A) $\pi/2$ (B) $\pi/4$ (C) $\pi^2/2$ (D) $\pi^2/4$

Sol: $\int_0^{\pi} \frac{x \tan x}{\sec x \cdot \operatorname{cosec} x} dx$

$$= \frac{\pi}{2} \int_0^{\pi} \frac{\tan x}{\sec x \cos x} dx$$

$$= \frac{\pi}{2} \int_0^{\pi} \sin^2 x dx$$

$$= \frac{\pi}{2} \int_0^{\pi} \frac{1 - \cos 2x}{2} dx$$

$$= \frac{\pi}{2} \left[\frac{x}{2} - \frac{1}{4} \sin 2x \right]_0^{\pi}$$

$$= \frac{\pi}{2} \left[\left(\frac{\pi}{2} - 0 \right) - (0 - 0) \right] = \frac{\pi^2}{4}$$

Ans: (D)

24. $\int \sqrt{5-2x+x^2} dx =$

(A) $\frac{x-1}{2} \sqrt{5+2x+x^2} + 2 \log \left| (x-1) + \sqrt{5+2x+x^2} \right| + C$

(B) $\frac{x-1}{2} \sqrt{5-2x+x^2} + 2 \log \left| (x+1) + \sqrt{x^2+2x+5} \right| + C$

(C) $\frac{x-1}{2} \sqrt{5-2x+x^2} + 2 \log \left| (x-1) + \sqrt{5-2x+x^2} \right| + C$

(D) $\frac{x}{2} \sqrt{5-2x+x^2} + 4 \log \left| (x+1) + \sqrt{x^2-2x+5} \right| + C$

Sol: $\int \sqrt{5-5x+x^2} dx = \int \sqrt{(x-1)^2 + 2^2} dx$

$$= \frac{x-1}{2} \sqrt{5-2x+x^2} + 2 \log \left| x-1 + \sqrt{5-2x+x^2} \right| + c$$

Ans: (C)

25. The area of the region bounded by the line $y = x+1$, and the lines $x = 3$ and $x = 5$ is

(A) $\frac{11}{2}$ sq. units

(B) 10 sq. units

(C) 7 sq. units

(D) $\frac{7}{2}$ sq. units

Sol: $A = \int_3^5 x+1 dx = \frac{x^2}{2} + x \Big|_3^5 = \frac{25}{2} - \frac{9}{2} + 5 - 3$

$$= \frac{16}{2} + 2 = 8 + 2 = 10$$

Ans: (B)

26. If a curve passes through the point (1,1) and at any point (x,y) on the curve, the product of the slope of its tangent and x co-ordinate of the point is equal to the y co-ordinate of the point, then the curve also passes through the point

(A) (-1,2)

(B) (2,2)

(C) $(\sqrt{3},0)$

(D) (3,0)

Sol: $x \frac{dy}{dx} = y$

$$\Rightarrow \frac{1}{y} dy = \frac{1}{x} dx$$

$$\Rightarrow \log y = \log x + C$$

(1, 1) lie on it

$$\Rightarrow \log 1 = \log 1 + C \Rightarrow C = 0$$

$\therefore \log y = \log x$

$\Rightarrow y = x$

Ans: (B)

27. The degree of the differential equation $1 + \left(\frac{dy}{dx}\right)^2 + \left(\frac{d^2y}{dx^2}\right)^2 = \sqrt[3]{\frac{d^2y}{dx^2} + 1}$ is

- (A) 1 (B) 6 (C) 2 (D) 3

Sol: $1 + (y')^2 + (y'')^2 = (y'' + 1)^{1/3}$

Cubing on B-S

$(1 + (y')^2 + (y'')^2)^3 = y'' + 1$

On verification degree = 6

\therefore (B) is the correct answer

Ans: (B)

28. If $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$ then

- (A) \vec{a} and \vec{b} are coincident. (B) \vec{a} and \vec{b} are perpendicular.
 (C) Inclined to each other at 60° . (D) \vec{a} and \vec{b} are parallel.

Sol: $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}| \Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = |\vec{a}|^2 + |\vec{b}|^2 - 2\vec{a} \cdot \vec{b}$

$4\vec{a} \cdot \vec{b} = 0$

$\vec{a} \cdot \vec{b} = 0$

$\vec{a} \perp \vec{b}$

Ans: (B)

29. The component of \hat{i} in the direction of the vector $\hat{i} + \hat{j} + 2\hat{k}$ is

- (A) $6\sqrt{6}$ (B) $\sqrt{6}$ (C) $\frac{\sqrt{6}}{6}$ (D) 6


Sol: $\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{1}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}} = \frac{\sqrt{6}}{6}$

Ans: (C)

30. In the interval $(0, \pi/2)$, area lying between the curves $y = \tan x$ and $y = \cot x$ and the X-axis is

- (A) $4\log 2$ sq. units (B) $3\log 2$ sq. units (C) $\log 2$ sq. units (D) $2\log 2$ sq. units

Sol:



$$A = 2 \int_0^{\frac{\pi}{4}} \tan x dx = 2 \log \sec x \Big|_0^{\pi/4} = 2 \log \sqrt{2} = \log 2$$

Ans: (C)

31. If $\vec{a} + 2\vec{b} + 3\vec{c} = \vec{0}$ and $(\vec{a} \times \vec{b}) + (\vec{b} \times \vec{c}) + (\vec{c} \times \vec{a}) = \lambda(\vec{b} \times \vec{c})$ then the value of λ is equal to

- (A) 4 (B) 2 (C) 6 (D) 3

Sol: $\vec{a} + 2\vec{b} + 3\vec{c} = \vec{0}$ [$\times \vec{b}$ both sides]

$$\vec{a} \times \vec{b} + 3\vec{c} \times \vec{b} = \vec{0}$$

$$\vec{a} \times \vec{b} = 3\vec{b} \times \vec{c}$$

$$\vec{a} + 2\vec{b} + 3\vec{c} = \vec{0} \quad (\vec{c} \times) \text{ on both sides}$$

$$\vec{c} \times \vec{a} + 2\vec{c} \times \vec{b} = \vec{0}$$

$$\vec{c} \times \vec{a} = 2\vec{b} \times \vec{c}$$

Ans: (C)

32. If a line makes an angle of $\frac{\pi}{3}$ with each X and Y axis then the acute angle made by Z -axis is

- (A) $\frac{\pi}{2}$ (B) $\frac{\pi}{6}$ (C) $\frac{\pi}{4}$ (D) $\frac{\pi}{3}$

Sol: $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$


$$\frac{1}{2} + \frac{1}{2} + \cos^2 \gamma = 1 \Rightarrow \gamma = 90^\circ$$

Ans: (A)

33. The length of perpendicular drawn from the point $(3, -1, 11)$ to the line $\frac{x}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ is

- (A) $\sqrt{33}$ (B) $\sqrt{66}$ (C) $\sqrt{53}$ (D) $\sqrt{29}$

Sol:



$$\vec{BA} = 3\hat{i} - 3\hat{j} + 8\hat{k}$$

$$\vec{BC} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$\vec{BA} \times \vec{BC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -3 & 8 \\ 2 & 3 & 4 \end{vmatrix} = -36\hat{i} + 4\hat{j} + 15\hat{k}$$

$$\vec{BC} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$\perp r \text{ distance} = \frac{\sqrt{1296+16+225}}{\sqrt{29}} = \sqrt{53}$$

Ans: (C)

34. The equation of the plane through the points $(2,1,0), (3,2,-2)$ and $(3,1,7)$ is

- (A) $6x-3y+2z-7=0$ (B) $3x-2y+6z-27=0$ (C) $7x-9y-z-5=0$ (D) $2x-3y+4z-27=0$

Sol:
$$\begin{vmatrix} x-2 & y-1 & z \\ 1 & 1 & -2 \\ 1 & 0 & 7 \end{vmatrix} = 0$$

$$\Rightarrow 7x-9y-z-5=0$$

Ans: (C)

35. The point of intersection of the line $x+1 = \frac{y+3}{3} = \frac{-z+2}{2}$ with the plane $3x+4y+5z=10$ is

- (A) $(2,6,-4)$ (B) $(-2,6,-4)$ (C) $(2,6,4)$ (D) $(2,-6,-4)$

Sol: Any point on the line $= (\lambda-1, 3\lambda-3, -2\lambda+2)$

This lies on plane

$$\therefore \lambda = 3$$


$$\therefore \text{Point } (2,6,-4)$$

Ans: (A)

36. If $(2,3,-1)$ is the foot of the perpendicular from $(4,2,1)$ to a plane, then the equation of the plane is

- (A) $2x-y+2z=0$ (B) $2x-y+2z+1=0$ (C) $2x+y+2z-5=0$ (D) $2x+y+2z-1=0$

Sol:



Equation of the plane is $2x-y+2z+k=0$ (1)

$(2,3,-1)$ lies on (1)

$$\therefore k = 1$$

Ans: (B)

37. $|\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = 144$ and $|\vec{a}| = 4$ then $|\vec{b}|$ is equal to

- (A) 8 (B) 12 (C) 4 (D) 3

Sol: $(|\vec{a}| |\vec{b}|)^2 = 144$

$$\Rightarrow |\vec{a}| |\vec{b}| = 12$$

$$\Rightarrow |\vec{b}| = \frac{12}{4} = 3$$

Ans: (D)

38. If A and B are events such that $P(A) = \frac{1}{4}$, $P(A/B) = \frac{1}{2}$ and $P(B/A) = \frac{2}{3}$ then $P(B)$ is

(A) $\frac{2}{3}$

(B) $\frac{1}{6}$

(C) $\frac{1}{2}$

(D) $\frac{1}{3}$

Sol: $P(A) \cdot P\left(\frac{B}{A}\right) = P(B) \cdot P\left(\frac{A}{B}\right)$

$$= \frac{1}{4} \times \frac{2}{3} = P(B) \cdot \frac{1}{2}$$

$$P(B) = \frac{1}{3}$$

Ans: (D)

39. A bag contains $2n+1$ coins. It is known that n of these coins have head on both sides whereas the other $n+1$ coins are fair. One coin is selected at random and tossed. If the probability that toss results in heads is $\frac{31}{42}$, then the value of n is

(A) 8

(B) 5

(C) 10

(D) 6

Sol: n coins - Two headers (TH)

$n+1$ coins - Fair

$$P(x) = \frac{n}{2n+1} \cdot 1 + \frac{n+1}{2n+1} \cdot \frac{1}{2} = \frac{31}{42} \quad \Rightarrow n=10$$

Ans: (C)

40. Let $A = \{x, y, z, u\}$ and $B = \{a, b\}$. A function $f : A \rightarrow B$ is selected randomly. The probability that the function is an onto function is

(A) $\frac{5}{8}$

(B) $\frac{7}{8}$

(C) $\frac{1}{35}$


(D) $\frac{1}{8}$

Sol: $n(s) = 2^4 = 16$

$$n(E) = 2^4 - 2 = 14 \quad P(E) = \frac{14}{16} = \frac{7}{8}$$

Ans: (B)

41. The shaded region in the figure given is the solution of which of the inequations?




(A) $x + y \geq 7, 2x - 3y + 6 \geq 0, x \geq 0, y \geq 0$

(B) $x + y \leq 7, 2x - 3y + 6 \geq 0, x \geq 0, y \geq 0$

(C) $x + y \leq 7, 2x - 3y + 6 \leq 0, x \geq 0, y \geq 0$

(D) $x + y \geq 7, 2x - 3y + 6 \leq 0, x \geq 0, y \geq 0$

Sol:



$2x - 3y + 6 \geq 0$

Ans: (B)

42. If $f(x) = ax + b$, where a and b are integers, $f(-1) = -5$ and $f(3) = 3$ then a and b are respectively

(A) 0, 2

(B) -3, -1

(C) 2, 3

(D) 2, -3

Sol: $f(-1) = -5$

$-a + b = -5 \dots (1)$

$(1) - (2) \quad -4a = -8$

$-2 + b = -5$

$b = -3$

$(a, b) = (2, -3)$

Ans: (D)

$f(3) = 3$

$3a + b = 3 \dots (2)$

$\Rightarrow a = 2$

43. The value of $e^{\log_{10} \tan 1^\circ + \log_{10} \tan 2^\circ + \log_{10} \tan 3^\circ + \dots + \log_{10} \tan 89^\circ}$ is

(A) $\frac{1}{e}$

(B) 0

(C) 1

(D) 3

Sol: $e^{\log_{10} \tan 1^\circ \cdot \cot 1^\circ + \dots + \log_{10} \tan 45^\circ} = e^{\log_{10} 1} = e^0 = 1$

Ans: (C)

44. The value of $\begin{vmatrix} \sin^2 14^\circ & \sin^2 66^\circ & \tan 135^\circ \\ \sin^2 66^\circ & \tan 135^\circ & \sin^2 14^\circ \\ \tan 135^\circ & \sin^2 14^\circ & \sin^2 66^\circ \end{vmatrix}$ is

(A) 1

(B) -1

(C) 2

(D) 0

Sol: WRONG QUESTION

Ans: ()

45. The modulus of the complex number $\frac{(1+i)^2(1+3i)}{(2-6i)(2-2i)}$ is

(A) $\frac{1}{\sqrt{2}}$

(B) $\frac{4}{\sqrt{2}}$

(C) $\frac{\sqrt{2}}{4}$

(D) $\frac{2}{\sqrt{2}}$

Sol: $\left| \frac{(1+i)^2(1+3i)}{(2-6i)(2-2i)} \right| = \frac{|(1+i)|^2 |1+3i|}{|2-6i| |2-2i|}$

$= \frac{(\sqrt{1+1})^2 \sqrt{1+9}}{\sqrt{4+36} \sqrt{4+4}} = \frac{2 \times \sqrt{10}}{\sqrt{40} 2\sqrt{2}} = \frac{1}{2\sqrt{2}} = \frac{\sqrt{2}}{4}$

Ans: (C)

46. Given that a, b and x are real numbers and $a < b, x < 0$ then

(A) $\frac{a}{x} < \frac{b}{x}$

(B) $\frac{a}{x} > \frac{b}{x}$

(C) $\frac{a}{x} \leq \frac{b}{x}$

(D) $\frac{a}{x} \geq \frac{b}{x}$

Sol: $a < b$

$4 > 5$

$x < 0$

$x = -2$

$\left. \begin{matrix} \frac{4}{-2} > \frac{5}{-2} \\ -2 > -2.5 \end{matrix} \right\} \frac{a}{x} > \frac{b}{x}$

Ans: (B)

47. Ten chairs are numbered as 1 to 10. Three women and two men wish to occupy one chair each. First the women choose the chairs marked 1 to 6, then the men choose the chairs from the remaining. The number of possible ways is

(A) ${}^6C_3 \times {}^4P_2$

(B) ${}^6C_3 \times {}^4C_2$

(C) ${}^6P_3 \times {}^4C_2$

(D) ${}^6P_3 \times {}^4P_2$

Sol: Women select 3 chairs in 6C_3 ways. Men select 2 chairs in 4C_2 ways

Total = ${}^6C_3 \times {}^4C_2$

Ans: (B)

48. Which of the following is an empty set?

(A) $\{x : x^2 - 9 = 0, x \in \mathbb{R}\}$ (B) $\{x : x^2 - 1 = 0, x \in \mathbb{R}\}$ (C) $\{x : x^2 = x + 2, x \in \mathbb{R}\}$ (D) $\{x : x^2 + 1 = 0, x \in \mathbb{R}\}$

Sol: $x^2 + 1 = 0, x \in \mathbb{R}$ is an empty set

Ans: (D)

49. n^{th} term of the series $1 + \frac{3}{7} + \frac{5}{7^2} + \frac{1}{7^2} + \dots$ is

(A) $\frac{2n-1}{7^n}$

(B) $\frac{2n-1}{7^{n-1}}$

(C) $\frac{2n+1}{7^{n-1}}$

(D) $\frac{2n+1}{7^n}$

Sol: $P(n): 1 + \frac{3}{7} + \frac{5}{7^2} + \frac{1}{7^2} + \dots$

$P(1): \frac{2 \cdot 1 - 1}{7^{1-1}} = 1$

$P(2): \frac{2 \times 2 - 1}{7^{2-1}} = \frac{3}{7}$

Ans: (B)

50. If $p\left(\frac{1}{q} + \frac{1}{r}\right), q\left(\frac{1}{r} + \frac{1}{p}\right), r\left(\frac{1}{p} + \frac{1}{q}\right)$ are in A.P., then p, q, r

(A) are in A.P.

(B) are not in A.P.

(C) are not in G.P.

(D) are in G.P.

Sol:

$p\left(\frac{1}{q} + \frac{1}{r} + \frac{1}{p} - \frac{1}{p}\right), q\left(\frac{1}{r} + \frac{1}{p} + \frac{1}{q} - \frac{1}{q}\right)$

$r\left(\frac{1}{p} + \frac{1}{q} + \frac{1}{r} - \frac{1}{r}\right)$

$pk - 1, qk - 1, rk - 1$

p, q, r are A.P

k is $\left(\frac{1}{p} + \frac{1}{q} + \frac{1}{r}\right)$ are in A.P

Ans: (A)

51. A line passes through (2,2) and is perpendicular to the line $3x + y = 3$. Its y -intercept is

(A) 1

(B) $\frac{1}{3}$

(C) $\frac{4}{3}$

(D) $\frac{2}{3}$

Sol: $3x + y = 3$

$y = -3x + 3$

$m_1 = 3$

$m_2 = \frac{1}{3}$

$y - 2 = \frac{1}{3}(x - 2)$

$x = 0$

$y - 2 = -\frac{2}{3}$

$y = 2 - \frac{2}{3} = \frac{4}{3}$

Ans: (C)

52. The distance between the foci of a hyperbola is 16 and its eccentricity is $\sqrt{2}$. Its equation is

- (A) $2x^2 - 3y^2 = 7$ (B) $x^2 - y^2 = 32$ (C) $y^2 - x^2 = 32$ (D) $\frac{x^2}{4} - \frac{y^2}{9} = 1$

Sol: $2ae = 16$

$$2a\sqrt{2} = 16$$

$$a = \frac{16}{2\sqrt{2}}$$

$$a = 4\sqrt{2}$$

$$x^2 - y^2 = 32$$

Ans: (B)

53. If $\lim_{x \rightarrow 0} \frac{\sin(2+x) - \sin(2-x)}{x} = A \cos B$, then the values of A and B respectively are

- (A) 2,1 (B) 2,2 (C) 1,1 (D) 1,2

Sol: $\lim_{x \rightarrow 0} 2 \cos[2] \frac{\sin x}{x}$

$$\lim_{x \rightarrow 0} 2 \cos 2 \cdot \frac{\sin x}{x}$$

$$2 \cos 2$$

$$A = 2, B = 2$$

Ans: (B)

54. If n is even and the middle term in the expansion of $\left(x^2 + \frac{1}{x}\right)^n$ is $924x^6$, then n is equal to

- (A) 12 (B) 10 (C) 8 (D) 14

Sol: Middle term $T_{\frac{n}{2}+1} = {}^n C_{\frac{n}{2}} \left(x^2\right)^{\frac{n}{2}} \left(\frac{1}{x}\right)^{\frac{n}{2}} = 924x^6$

$${}^n C_{\frac{n}{2}} x^2 = 924x^6$$

$$\frac{n}{2} = 6 \Rightarrow n = 12$$

Ans: (A)

55. The mean of 100 observations is 50 and their standard deviation is 5. Then the sum of squares of all observations is

- (A) 250000 (B) 50000 (C) 255000 (D) 252500

Sol: $\frac{\sum x_i}{100} = 50; \sigma = 5$

$$\frac{\sum x_i^2}{100} - \left(\frac{\sum x_i}{100}\right)^2 = 25$$

$$\frac{\sum x_i^2}{100} = 2500 + 25$$

$$\Rightarrow \sum x_i^2 = 252500$$

Ans: (D)

56. $f : R \rightarrow R$ and $g : [0, \infty) \rightarrow R$ are defined by $f(x) = x^2$ and $g(x) = \sqrt{x}$. Which one of the following is not true ?

- (A) $(f \circ g)(2) = 2$ (B) $(g \circ f)(4) = 4$ (C) $(g \circ f)(-2) = 2$ (D) $(f \circ g)(-4) = 4$

Sol: $f(g(2)) = (g(2))^2 = (\sqrt{2})^2 = 2 \rightarrow \text{True}$

$g(f(4)) = \sqrt{f(4)} = \sqrt{4^2} = 4 \rightarrow \text{True}$

$g(f(-2)) = \sqrt{f(-2)} = \sqrt{(-2)^2} = 2 \rightarrow \text{True}$

$f(g(-4)) = (g(-4))^2 = (\sqrt{-4})^2 \neq 4 \rightarrow \text{False}$

Ans: (D)

57. Let $f : R \rightarrow R$ be defined by $f(x) = 3x^2 - 5$ and $g : R \rightarrow R$ by $g(x) = \frac{x}{x^2 + 1}$ then $g \circ f$ is

- (A) $\frac{3x^2}{x^4 + 2x^2 - 4}$ (B) $\frac{3x^2 - 5}{9x^4 - 30x^2 + 26}$ (C) $\frac{3x^2}{9x^4 + 30x^2 - 2}$ (D) $\frac{3x^2 - 5}{9x^4 - 6x^2 + 26}$

Sol: $g(f(x)) = \frac{3x^2 - 5}{(3x^2 - 5)^2 + 1} = \frac{3x^2 - 5}{9x^4 - 30x^2 + 26}$

Ans: (B)

58. Let the relation R be defined in N by aRb if $3a + 2b = 27$ then R is

- (A) $\{(1,12), (3,9), (5,6), (7,3), (9,0)\}$ (B) $\{(1,12), (3,9), (5,6), (7,3)\}$
 (C) $\{(2,1), (9,3), (6,5), (3,7)\}$ (D) $\left\{\left(0, \frac{27}{2}\right), (1,12), (3,9), (5,6), (7,3)\right\}$

Sol: $a \in N, b \in N$

From option B all satisfies condition

Ans: (B)

59. Let $f(x) = \sin 2x + \cos 2x$ and $g(x) = x^2 - 1$, then $g(f(x))$ is invertible in the domain

- (A) $x \in \left[\frac{-\pi}{2}, \frac{\pi}{2}\right]$ (B) $x \in \left[\frac{-\pi}{4}, \frac{\pi}{4}\right]$ (C) $x \in \left[0, \frac{\pi}{4}\right]$ (D) $x \in \left[\frac{-\pi}{8}, \frac{\pi}{8}\right]$

Sol: $g(f(x))$

$$g\left[\left(\frac{1}{\sqrt{2}} \sin 2x + \frac{1}{\sqrt{2}} \cos 2x\right) \sqrt{2}\right]$$

$$g\left[\sqrt{2} \sin\left(2x + \frac{\pi}{4}\right)\right]$$

$$g(f(x)) = 2 \sin^2 \left(2x + \frac{\pi}{4} \right) - 1$$

$$= -\cos \left[4x + \frac{\pi}{2} \right]$$

$g(f(x))$ is invertible

$$0 \leq 4x + \frac{\pi}{2} \leq \pi$$

$$-\frac{\pi}{8} \leq x \leq \frac{\pi}{8}$$

Ans: (D)

60. The contrapositive of the statement, "if two lines do not intersect in the same plane, then they are parallel" is

- (A) If two lines are not parallel then they do not intersect in the same plane.
- (B) If two lines are not parallel then they intersect in the same plane.
- (C) If two lines are parallel then they do not intersect in the same plane.
- (D) If two lines are parallel then they intersect in the same plane.

Sol: p : Two lines do not intersect in same plane

Q : Two lines are parallel

Contrapositive: $\sim q \rightarrow \sim p$

If two lines not parallel they intersect in same plane

Ans: (B)

Key Answers:

1. B	2. B	3. B	4. B	5. C	6. B	7. C	8. C	9. B	10. B
11. C	12. D	13. B	14. C	15. A	16. D	17. C	18. C	19. A	20. B
21. A	22. A	23. D	24. C	25. B	26. B	27. B	28. B	29. C	30. C
31. C	32. A	33. C	34. C	35. A	36. B	37. D	38. D	39. C	40. B
41. B	42. D	43. C	44.	45. C	46. B	47. B	48. D	49. B	50. A
51. C	52. B	53. B	54. A	55. D	56. D	57. B	58. B	59. D	60. B

44. (Wrong Question)